

Detection of the Neutrino Fluxes from Several Sources

D.L. Khokhlov
*Sumy State University**
 (Dated: February 2, 2008)

It is considered the detection of neutrinos moving from the opposite directions. The states of the particle of the detector interacting with the neutrinos are connected with the P-transformation. Hence only a half of neutrinos gives contribution into the superposition of the neutrino states. Taking into account the effect of the opposite neutrino directions the total neutrino flux from several sources are in the range 0.5–1 of that without the effect. The neutrino flux from nuclear reactors measured in the KamLAND experiment is $0.611 \pm 0.085(\text{stat}) \pm 0.041(\text{syst})$ of the expected flux. Calculations for the conditions of the KamLAND experiment yield the neutrino flux taking into account the effect of the opposite neutrino directions, 0.555, of that without the effect that may account for the neutrino flux observed in the KamLAND experiment.

PACS numbers: 03.65.-w, 28.50.Hw

Let two equal neutrino fluxes move towards the neutrino detector from the opposite directions. The first neutrino flux moves from the x direction, and the second neutrino flux moves from the $-x$ direction. The probability of the weak interaction of the particle of the detector with a single neutrino is given by

$$w = |\langle \psi_1 \psi_2 | g | \psi_\nu \psi_p \rangle|^2 \quad (1)$$

where g is the weak coupling, ψ_p is the state of the particle of the detector, ψ_ν is the state of neutrino, ψ_1, ψ_2 are the states of the particles arising in the interaction. The probability given by eq. (1) is too small to register the event in a single interaction of the particle of the detector with the neutrino. The probability of the event is a sum of probabilities of a number of interactions.

To define the probability of interaction of the particle of the detector with two neutrinos moving from the opposite directions one should form the superposition of the particle of the detector and neutrino states. The state of the particle interacting with the neutrino moving from the x direction and the state of the particle interacting with the neutrino moving from the $-x$ direction are connected with the P-transformation

$$|\psi_p(-x)\rangle = P|\psi_p(x)\rangle. \quad (2)$$

Then the superposition of the particle and neutrino states should be given by

$$\begin{aligned} |\psi\rangle &= \frac{1}{\sqrt{2}} |\psi_\nu\rangle [|\psi_p(x)\rangle + |\psi_p(-x)\rangle] \\ &= \frac{1}{\sqrt{2}} |\psi_\nu\rangle [|\psi_p(x)\rangle + P|\psi_p(x)\rangle] \\ &= \frac{1}{\sqrt{2}} [|\psi_\nu\rangle + P|\psi_\nu\rangle] |\psi_p(x)\rangle. \end{aligned} \quad (3)$$

Since the P-transformation is forbidden for neutrino, one cannot form the superposition given by eq. (3). Hence only a half of neutrinos moving from the opposite directions gives contribution into the superposition of the neutrino states. Therefore the sum of probabilities of interaction of the particle with neutrinos moving from the opposite directions is two times smaller than that for the case when all neutrinos move in one and the same direction

$$w_{\uparrow\downarrow} = 0.5 w_{\uparrow\uparrow} \quad (4)$$

Let the detector measure the neutrino fluxes F_i from N sources placed in the plane xy . When not taking into account the effect of the opposite neutrino directions given by eq. (4) the total neutrino flux is

$$F = \sum_{i=1}^N F_i. \quad (5)$$

To take into account the effect of the opposite neutrino directions given by eq. (4) write down the neutrino flux in the form

$$F_i = F_{ix} + F_{iy} \quad (6)$$

where

$$F_{ix} = F_i \cos^2 \alpha \quad (7)$$

$$F_{iy} = F_i \sin^2 \alpha \quad (8)$$

where α is the angle between the direction to the source and the axis x . When taking into account the effect of the opposite neutrino directions given by eq. (4) the total neutrino flux is

$$F^* = 0.5 \left(\sum_{i=1}^N F_i + \sum_{i=1}^N \vec{F}_{ix} + \sum_{i=1}^N \vec{F}_{iy} \right) \quad (9)$$

*Electronic address: khokhlov@cafe.sumy.ua

where one should sum the fluxes $\vec{F}_{ix}$, $\vec{F}_{iy}$ as vectors. One can see that the neutrino flux given by eq. (9) are within the range 0.5–1 of the neutrino flux given by eq. (5).

In the KamLAND experiment [1], it is measured the flux of $\bar{\nu}_e$'s from distant nuclear reactors. The ratio of the number of observed events to the expected number of events is reported [2] to be $0.611 \pm 0.085(\text{stat}) \pm 0.041(\text{syst})$. Using the data on the location and thermal power of reactors [3] one can estimate the neutrino flux from reactors which is approximately proportional to the thermal power flux $P_{th}/4\pi L^2$, where P_{th} is the reactor thermal power, L is the distance from the detector to the reactor. Hence one can estimate the effect of the opposite neutrino directions through the ratio of the neutrino flux calculated with eq. (9) and that calculated with

eq. (5). More than 99% of the neutrino flux measured in the KamLAND experiment comes from Japanese and Korean nuclear reactors. So in the first approximation one can calculate the neutrino flux only from Japanese and Korean nuclear reactors. Calculations for the conditions of the KamLAND experiment yield the ratio of the neutrino flux taking into account the effect of the opposite neutrino directions and that without this effect, 0.555. Thus the effect of the opposite neutrino directions may account for the neutrino flux observed in the KamLAND experiment. Note that the value 0.555 is obtained with the use of the specification data on the reactor thermal power. To obtain the real value one should use the operation data on the reactor thermal power for the period of measurement.

[1] <http://www.awa.tohoku.ac.jp/KamLAND/>

[2] K. Eguchi *et al.*, KamLAND Collaboration, submitted to Phys. Rev. Lett., hep-ex/0212021.

[3] <http://www.insc.anl.gov/>